


5  
SCI08  
Dup

# SCIENCE

20 OCTOBER 1995  
Vol. 270 • PAGES 349-530




## GENOME ISSUE

BEST AVAILABLE COPY

88XNCNG \*\*\*\*\* 3-DIGIT 530  
80258222810128AS 12/01/95 E 5235 74  
ELDO C KOENIG  
35005 M FAIRVIEW RD  
OCNOMOC MI 53066-3312

The Genome Project adds a new dimension to questions of gene expression in humans and model systems. A report on page 415 summarizes progress in the *renobabdiis elegans* Genome Project and indicates the ways information about sequences can be used.

News stories, Articles, Perspectives, Policy Forums, and Reports focus on technological developments, clinical applications, and ethical concerns resulting from the burgeoning of genomic information. [*C. elegans* image: F. Maduro and D. Pilgrim, University of Alberta]



## REPORTS

Seismogenic Ages for Earthquake  
 recurrence Intervals and Debris Flow Fan Depo-  
 sition, Owens Valley, California  
 R. Bierman, A. R. Gillespie, M. W. Caffee

Photoautotrophic Microbial  
 ecosystems in Deep Basalt Aquifers  
 O. Stevens and J. P. McKinley

Large Arctic Temperature Change at  
 Wisconsin-Holocene Glacial Transition  
 M. Cuffey, G. D. Clow, R. B. Alley, M. Stuiver,  
 D. Waddington, R. W. Saltus

Permeability in Earth's Lower Mantle:  
 Evidence from Seismic Anisotropy and  
 Rock Physics  
 J. Karato, S. Zhang, H.-R. Wenk

Large-Scale Interplanetary Magnetic  
 Field Configuration Revealed by Solar  
 Radio Bursts  
 J. J. Reiner, J. Fainberg, R. G. Stone

Role of Yeast Insulin-Degrading Enzyme  
 Homologs in Pheromone Processing and  
 Bud Site Selection  
 N. Adames, K. Blundell, M. N. Ashby, C.  
 Boone

Quantitative Monitoring of Gene  
 Expression Patterns with a  
 Complementary DNA Microarray  
 M. Schena, D. Shalon, R. W. Davis, P. O.  
 Brown

Gene Therapy in Peripheral Blood  
 Lymphocytes and Bone Marrow for  
 ADA<sup>-</sup> Immunodeficient Patients  
 C. Bordignon, L. D. Notarangelo, N. Nobili, G.  
 Ferrari, G. Casorati, P. Panina, E. Mazzolari, D.  
 Maggioni, C. Rossi, P. Servida, A. G. Ugazio, F.  
 Mavilio


T Lymphocyte-Directed Gene  
 Therapy for ADA<sup>-</sup> SCID: Initial  
 Trial Results After 4 Years  
 R. M. Blaese, K. W. Culver, A. D. Miller, C. S.  
 Carter, T. Fleisher, M. Clerici, G. Shearer, L.  
 Chang, Y. Chiang, P. Tolstoshev, J. J. Greenblatt,  
 S. A. Rosenberg, H. Klein, M. Berger, C. A.  
 Mullen, W. J. Ramsey, L. Muul, R. A. Morgan,  
 W. F. Anderson

Physical Map and Organization of  
 *Arabidopsis thaliana* Chromosome 4  
 R. Schmidt, J. West, K. Love,  
 Z. Lenehan, C. Lister, H. Thompson,  
 D. Bouchet, C. Dean

Serial Analysis of Gene Expression  
 V. E. Velculescu, L. Zhang,  
 B. Vogelstein, K. W. Kinzler

## TECHNICAL COMMENTS

The Radius of Gyration of an  
 Apomyoglobin Folding Intermediate  
 D. Eliezer, P. A. Jennings, P. E. Wright, S. Doniach,  
 K. O. Hodgson, H. Tsuruta



397

Good things in  
 small genomes

NOT AVAILABLE COPY

## AAAS Board of Directors

David J. Ayala  
 Incoming President  
 Herman  
 R. Colwell  
 President  
 Lubchenco  
 President-elect

Anne C. Roosevelt  
 Alan Schnesheim  
 Jean E. Taylor  
 Chang-Lin Tian  
 Nancy S. Weller

William T. Golden  
 Treasurer  
 Richard S. Nicholson  
 Executive Officer

SCIENCE (ISSN 0036-8075) is published weekly on Friday, except the last week in December, by the American Association for the Advancement of Science, 1333 H Street, NW, Washington, DC 20005. Second-class postage (publication No. 484460) paid at Washington, DC, and additional mailing offices. Copyright © 1995 by the American Association for the Advancement of Science. The title SCIENCE is a registered trademark of the AAAS. Domestic individual membership and subscription (51 issues): \$97 (\$50 allocated to subscription). Domestic institutional subscription (51 issues): \$228. Foreign postage extra: Mexico, Caribbean (surface mail) \$53; other countries (air assist delivery) \$93. First class, airmail, student and emeritus rates on request. Canadian rates with GST available upon request. GST #1254 88122. Printed in the U.S.A.

Change of address: allow 4 weeks, giving old and new addresses and 8-digit account number. Postmaster: Send change of address to Science, P.O. Box 1811, Danbury, CT 06813-1811. Single copy sales: \$7.00 per issue prepaid includes surface postage; bulk rates on request. Authorization to photocopy items for internal or personal use under circumstances not falling within the fair use provisions of the Copyright Act is granted by AAAS to libraries and other users registered with the Copyright Clearance Center (CCC) Transactional Reporting Service, provided that \$3.00 per article is paid directly to CCC, 27 Congress Street, Salem, MA 01970. The identification code for Science is 0036-8075/95 \$3.00. Science is indexed in the Reader's Guide to Periodical Literature and in several specialized indexes.

Ad1p sequence following Ser<sup>208</sup> and occurs within the domain of Ad1p that shows homology with hIDE (14). To delete the complete STE23 sequence and create the ste23Δ::URA3 mutation, polymerase chain reaction (PCR) primers (5'-TCGGAAGACCTCAT-TCTTGCTCATTTTGATATTGCTC- TGTAGATTG-TACTGAGAGTGAC-3'; and 5'-GCTACAAACAGC-GTGCAGCTTGAATGCCCGACATCTTCGACTGT-GCGGTATTTCACACCG-3') were used to amplify the URA3 sequence of pRS316, and the reaction product was transformed into yeast for one-step gene replacement [R. Rothstein, *Methods Enzymol.* 194, 281 (1991)]. To create the axl1Δ::LEU2 mutation contained on p114, a 5.0-kb Sal I fragment from pAXL1 was cloned into pUC19, and an internal 4.0-kb Hpa I-Xho I fragment was replaced with a LEU2 fragment. To construct the ste23Δ::LEU2 allele (a deletion corresponding to 931 amino acids) carried on p153, a LEU2 fragment was used to replace the 2.8-kb Pml I-Eco136 II fragment of STE23, which occurs within a 6.2-kb Hind III-Bgl II genomic fragment carried on pSP72 (Promega). To create YEpMFA1, a 1.6-kb Bam HI fragment containing MFA1, from pK16 [K. Kuchler, R. E. Sterne, J. Thome, *EMBO J.* 8, 3973 (1989)], was ligated into the Bam HI site of YEp351 [J. E. Hill, A. M. Myers, T. J. Koerner, A. Tzagoloff, *Yeast* 2, 163 (1986)].

4. J. Chant and I. Herskowitz, *Cell* 65, 1203 (1991).

5. B. W. Matthews, *Acc. Chem. Res.* 21, 333 (1988).

6. K. Kuchler, H. G. Dohlman, J. Thome, *J. Cell Biol.* 120, 1203 (1993); R. Koling and C. P. Hollenberg, *EMBO J.* 13, 3261 (1994); C. Berkower, D. Loayza, S. Michaels, *Mol. Biol. Cell* 5, 1185 (1994).

7. A. Bender and J. R. Pringle, *Proc. Natl. Acad. Sci. U.S.A.* 86, 9976 (1989); J. Chant, K. Corrado, J. R. Pringle, I. Herskowitz, *Cell* 65, 1213 (1991); S. Powers, E. Gonzales, T. Christensen, J. Cubert, D. Broek, *ibid.*, p. 1225; H. O. Park, J. Chant, I. Herskowitz, *Nature* 365, 269 (1993); J. Chant, *Trends Genet.* 10, 328 (1994); — and J. R. Pringle, *J. Cell Biol.* 129, 751 (1995); J. Chant, M. Mischke, E. Mitchell, I. Herskowitz, J. R. Pringle, *ibid.*, p. 767.

8. G. F. Sprague Jr., *Methods. Enzymol.* 194, 77 (1991).

9. Single-letter abbreviations for the amino acid residues are as follows: A, Ala; C, Cys; D, Asp; E, Glu; F, Phe; G, Gly; H, His; I, Ile; K, Lys; L, Leu; M, Met; N, Asn; P, Pro; Q, Gln; R, Arg; S, Ser; T, Thr; V, Val; W, Trp; and Y, Tyr.

10. A W303 1A derivative, SY2625 (MATa ura3-1 leu2-3, 112 trp1-1 ade2-1 can1-100 sst1Δ mfa2Δ::FUS1- lacZ his3Δ::FUS1-HIS3), was the parent strain for the mutant search. SY2625 derivatives for the mating assays, secreted pheromone assays, and the pulse-chase experiments included the following strains: Y49 (ste22-1), Y115 (mfa1Δ::LEU2), Y142 (axl1::URA3), Y173 (axl1Δ::LEU2), Y220 (axl1::URA3 ste23Δ::URA3), Y221 (ste23Δ::URA3), Y231 (axl1Δ::LEU2 ste23Δ::LEU2), and Y233 (ste23Δ::LEU2). MATa derivatives of SY2625 included the following strains: Y199 (SY2625 made MATa), Y278 (ste22-1), Y195 (mfa1Δ::LEU2), Y196 (axl1Δ::LEU2), and Y197 (axl1::URA3). The EG123 (MATa leu2 ura3 trp1 can1 his4) genetic background was used to create a set of strains for analysis of bud site selection. EG123 derivatives included the following strains: Y175 (axl1Δ::LEU2), Y223 (axl1::URA3), Y234 (ste23Δ::LEU2), and Y272 (axl1Δ::LEU2 ste23Δ::LEU2). MATa derivatives of EG123 included the following strains: Y214 (EG123 made MATa) and Y293 (axl1Δ::LEU2). All strains were generated by means of standard genetic or molecular methods involving the appropriate constructs (23). In particular, the axl1 ste23 double mutant strains were created by crossing of the appropriate MATa ste23 and MATa axl1 mutants, followed by sporulation of the resultant diploid and isolation of the double mutant from nonparental di-type tetrads. Gene disruptions were confirmed with either PCR or Southern (DNA) analysis.

p129 is a YEp352 [J. E. Hill, A. M. Myers, T. J. Koerner, A. Tzagoloff, *Yeast* 2, 163 (1986)] plasmid containing a 5.5-kb Sal I fragment of pAXL1. p151 was derived from p129 by insertion of a linker at the Bgl II site within AXL1, which led to an in-frame insertion of the hemagglutinin (HA) epitope (DOYPPYDVPDYA) (29) between amino acids 854 and 855 of the AXL1 prod-

uct. pC225 is a KS+ (Stratagene) plasmid containing a 0.5-kb Bam HI-Sst I fragment from pAXL1. Substitution mutations of the proposed active site of Axl1p were created with the use of pC225 and site-specific mutagenesis involving appropriate synthetic oligonucleotides (axl1-H68A, 5'-GTGCTCACAAAGCGCT-GCCAAACCGGC-3'; axl1-E71A, 5'-AAGAATCAT-GTGCGCACAAAGGTGCGC-3'; and axl1-E71D, 5'-AAGAATCATGTGATCACAAAGGTGCGC-3'). The mutations were confirmed by sequence analysis. After mutagenesis, the 0.4-kb Bam HI-Msc I fragment from the mutagenized pC225 plasmids was transferred into pAXL1 to create a set of pRS316 plasmids carrying different AXL1 alleles, p124 (axl1-H68A), p130 (axl1-E71A), and p132 (axl1-E71D). Similarly, a set of HA-tagged alleles carried on YEp352 were created after replacement of the p151 Bam HI-Msc I fragment, to generate p161 (axl1-E71A), p162 (axl1-

H68A), and p163 (axl1-E71D).

32. We thank J. Becker and S. Michaels for providing a-factor antibodies; S. Michaels for discussing unpublished results and helping with the pulse-chase experiments; J. Brown, J. Chant, and S. Sanoers for their input concerning bud site selection experiments; M. Raymond, F. Tamino, and M. Whiteway for plasmids; M. Marra for providing the STE23 genomic fragment; and H. Bussey, J. Brown, N. Davis, T. Favero, C. de Hoog, and S. Kim for comments on the manuscript. Supported by a grant to C.B. from the Natural Sciences and Engineering Research Council of Canada. Support for M.N.A. was from a California Tobacco-Related Disease Research Program postdoctoral fellowship (4FT-0083).

22 June 1995; accepted 21 August 1995

## Quantitative Monitoring of Gene Expression Patterns with a Complementary DNA Microarray

Mark Schena,\* Dari Shalon,† Ronald W. Davis, Patrick O. Brown‡

A high-capacity system was developed to monitor the expression of many genes in parallel. Microarrays prepared by high-speed robotic printing of complementary DNAs on glass were used for quantitative expression measurements of the corresponding genes. Because of the small format and high density of the arrays, hybridization volumes of 2 microliters could be used that enabled detection of rare transcripts in probe mixtures derived from 2 micrograms of total cellular messenger RNA. Differential expression measurements of 45 *Arabidopsis* genes were made by means of simultaneous, two-color fluorescence hybridization.

The temporal, developmental, topographical, histological, and physiological patterns in which a gene is expressed provide clues to its biological role. The large and expanding database of complementary DNA (cDNA) sequences from many organisms (1) presents the opportunity of defining these patterns at the level of the whole genome.

For these studies, we used the small flowering plant *Arabidopsis thaliana* as a model organism. *Arabidopsis* possesses many advantages for gene expression analysis, including the fact that it has the smallest genome of any higher eukaryote examined to date (2). Forty-five cloned *Arabidopsis* cDNAs (Table 1), including 14 complete sequences and 31 expressed sequence tags (ESTs), were used as gene-specific targets. We obtained the ESTs by selecting cDNA clones at random from an *Arabidopsis* cDNA library. Sequence analysis revealed that 28 of the 31 ESTs matched sequences

in the database (Table 1). Three additional cDNAs from other organisms served as controls in the experiments.

The 48 cDNAs, averaging ~1.0 kb, were amplified with the polymerase chain reaction (PCR) and deposited into individual wells of a 96-well microtiter plate. Each sample was duplicated in two adjacent wells to allow the reproducibility of the arraying and hybridization process to be tested. Samples from the microtiter plate were printed onto glass microscope slides in an area measuring 3.5 mm by 5.5 mm with the use of a high-speed arraying machine (3). The arrays were processed by chemical and heat treatment to attach the DNA sequences to the glass surface and denature them (3). Three arrays, printed in a single lot, were used for the experiments here. A single microtiter plate of PCR products provides sufficient material to print at least 500 arrays.

Fluorescent probes were prepared from total *Arabidopsis* mRNA (4) by a single round of reverse transcription (5). The *Arabidopsis* mRNA was supplemented with human acetylcholine receptor (AChR) mRNA at a dilution of 1:10,000 (w/w) before cDNA synthesis, to provide an internal standard for calibration (5). The resulting fluorescently labeled cDNA mixture was hybridized to an array at high stringency (6) and scanned

M. Schena and R. W. Davis, Department of Biochemistry, Beckman Center, Stanford University Medical Center, Stanford, CA 94305, USA.

D. Shalon and P. O. Brown, Department of Biochemistry and Howard Hughes Medical Institute, Beckman Center, Stanford University Medical Center, Stanford, CA 94305, USA.

\*These authors contributed equally to this work.

†Present address: Syntex, Palo Alto, CA 94303, USA.

‡To whom correspondence should be addressed. E-mail: pbrown@cirmg.stanford.edu

BEST AVAILABLE COPY


with a laser (3). A high-sensitivity scan gave signals that saturated the detector at nearly all of the *Arabidopsis* target sites (Fig. 1A). Calibration relative to the AChR mRNA standard (Fig. 1A) established a sensitivity limit of  $\sim 1:50,000$ . No detectable hybridization was observed to either the rat glucocorticoid receptor (Fig. 1A) or the yeast TRP4 (Fig. 1A) targets even at the highest scanning sensitivity. A moderate-sensitivity scan

of the same array allowed linear detection of the more abundant transcripts (Fig. 1B). Quantitation of both scans revealed a range of expression levels spanning three orders of magnitude for the 45 genes tested (Table 2). RNA blots (7) for several genes (Fig. 2) corroborated the expression levels measured with the microarray to within a factor of 5 (Table 2).

Differential gene expression was investi-

gated with a simultaneous, two-color hybridization scheme, which served to minimize experimental variation inherent in the comparison of independent hybridizations. Fluorescent probes were prepared from two mRNA sources with the use of reverse transcriptase in the presence of fluorescein- and lissamine-labeled nucleotide analogs, respectively (5). The two probes were then mixed together in equal proportions, hybridized to a single array, and scanned separately for fluorescein and lissamine emission after independent excitation of the two fluorophores (3).

To test whether overexpression of a single gene could be detected in a pool of total *Arabidopsis* mRNA, we used a microarray to analyze a transgenic line overexpressing the single transcription factor *HAT4* (8). Fluorescent probes representing mRNA from wild-type and *HAT4*-transgenic plants were labeled with fluorescein and lissamine, respectively; the two probes were then mixed and hybridized to a single array. An intense hybridization signal was observed at the position of the *HAT4* cDNA in the lissamine-specific scan (Fig. 1D), but not in the fluorescein-specific scan of the same array (Fig. 1C). Calibration with AChR mRNA added to the fluorescein and lissamine cDNA synthesis reactions at dilutions of 1:10,000 (Fig. 1C) and 1:100 (Fig. 1D), respectively, revealed a 50-fold elevation of *HAT4* mRNA in the transgenic line relative to its abundance in wild-type plants (Table 2). This magnitude of *HAT4* overexpression matched that inferred from the Northern (RNA) analysis within a factor of 2 (Fig. 2 and Table 2). Expression of all the other genes monitored on the array differed by less than a factor of 5 between *HAT4*-transgenic and wild-type plants (Fig. 1, C



1. Gene expression monitored with the use of cDNA microarrays. Fluorescent scans represented in pseudocolor correspond to hybridization intensities. Color bars were calibrated from the signal obtained from the use of known concentrations of human AChR mRNA in independent experiments. Numbers and letters on the axes mark the position of each cDNA. (A) High-sensitivity fluorescein scan after hybridization with a 1:1 mixture of fluorescein-labeled cDNA derived from wild-type plants. (B) Same array as in (A) but scanned at moderate sensitivity. (C and D) A single array was probed with a 1:1 mixture of fluorescein-labeled cDNA from wild-type plants and lissamine-labeled cDNA from *HAT4*-transgenic plants. The single array was scanned successively to detect the fluorescein fluorescence corresponding to mRNA from wild-type plants (C) and the lissamine fluorescence corresponding to mRNA from *HAT4*-transgenic plants (D). (E and F) A single array was probed with a 1:1 mixture of fluorescein-labeled cDNA from root tissue and lissamine-labeled cDNA from leaf tissue. The single array was then scanned successively to detect the fluorescein fluorescence corresponding to mRNAs expressed in roots (E) and the lissamine fluorescence corresponding to mRNAs expressed in leaves (F).




Fig. 2. Gene expression monitored with RNA (Northern) blot analysis. Designated amounts of mRNA from wild-type and *HAT4*-transgenic plants were spotted onto nylon membranes and probed with the cDNAs indicated. Purified human AChR mRNA was used for calibration.

and D, and Table 2). Hybridization of fluorescein-labeled glucocorticoid receptor cDNA (Fig. 1C) and lissamine-labeled TRP4 cDNA (Fig. 1D) verified the presence of the negative control targets and the lack of optical cross talk between the two fluorophores.

To explore a more complex alteration in expression patterns, we performed a second two-color hybridization experiment with fluorescein- and lissamine-labeled probes prepared from root and leaf mRNA, respectively. The scanning sensitivities for the two fluorophores were normalized by matching the signals resulting from AChR

mRNA, which was added to both cDNA synthesis reactions at a dilution of 1:1000 (Fig. 1, E and F). A comparison of the scans revealed widespread differences in gene expression between root and leaf tissue (Fig. 1, E and F). The mRNA from the light-regulated *CAB1* gene was ~500-fold more abundant in leaf (Fig. 1F) than in root tissue (Fig. 1E). The expression of 26 other genes differed between root and leaf tissue by more than a factor of 5 (Fig. 1, E and F).

The *HAT4*-transgenic line we examined has elongated hypocotyls, early flowering, poor germination, and altered pigmentation (8). Although changes in expression were

observed for *HAT4*, large changes in expression were not observed for any of the other 44 genes we examined. This was somewhat surprising, particularly because comparative analysis of leaf and root tissue identified 27 differentially expressed genes. Analysis of an expanded set of genes may be required to identify genes whose expression changes upon *HAT4* overexpression; alternatively, a comparison of mRNA populations from specific tissues of wild-type and *HAT4*-transgenic plants may allow identification of downstream genes.

At the current density of robotic printing, it is feasible to scale up the fabrication process to produce arrays containing 20,000 cDNA targets. At this density, a single array would be sufficient to provide gene-specific targets encompassing nearly the entire repertoire of expressed genes in the *Arabidopsis* genome (2). The availability of 20,274 ESTs from *Arabidopsis* (1, 9) would provide a rich source of templates for such studies.

The estimated 100,000 genes in the human genome (10) exceeds the number of *Arabidopsis* genes by a factor of 5 (2). This modest increase in complexity suggests that similar cDNA microarrays, prepared from the rapidly growing repertoire of human ESTs (1), could be used to determine the expression patterns of tens of thousands of human genes in diverse cell types. Coupling an amplification strategy to the reverse transcription reaction (11) could make it feasible to monitor expression even in minute tissue samples. A wide variety of acute and chronic physiological and pathological conditions might lead to characteristic changes in the patterns of gene expression in peripheral blood cells or other easily sampled tissues. In concert with cDNA microarrays for monitoring complex expression patterns, these tissues might therefore serve as sensitive *in vivo* sensors for clinical diagnosis. Microarrays of cDNAs could thus provide a useful link between human gene sequences and clinical medicine.

**Table 2.** Gene expression monitoring by microarray and RNA blot analyses; tg, *HAT4*-transgenic. See Table 1 for additional gene information. Expression levels (w/w) were calibrated with the use of known amounts of human AChR mRNA. Values for the microarray were determined from microarray scans (Fig. 1); values for the RNA blot were determined from RNA blots (Fig. 2).

Gene	Expression level (w/w)	
	Microarray	RNA blot
<i>CAB1</i>	1:48	1:83
<i>CAB1</i> (tg)	1:120	1:150
<i>HAT4</i>	1:8300	1:6300
<i>HAT4</i> (tg)	1:150	1:210
<i>ROC1</i>	1:1200	1:1800
<i>ROC1</i> (tg)	1:260	1:1300

**Table 1.** Sequences contained on the cDNA microarray. Shown is the position, the known or putative function, and the accession number of each cDNA in the microarray (Fig. 1). All but three of the ESTs used in this study matched a sequence in the database. NADH, reduced form of nicotinamide adenine dinucleotide; ATPase, adenosine triphosphatase; GTP, guanosine triphosphate.

Position	cDNA	Function	Accession number
a1, 2	AChR	Human AChR	-
a3, 4	EST3	Actin	H36236
a5, 6	EST6	NADH dehydrogenase	Z27010
a7, 8	AAC1	Actin 1	M20016
a9, 10	EST12	Unknown	U36594†
a11, 12	EST13	Actin	T45783
b1, 2	<i>CAB1</i>	Chlorophyll a/b binding	M85150
b3, 4	EST17	Phosphoglycerate kinase	T44490
b5, 6	G44	Gibberellic acid biosynthesis	L37126
b7, 8	EST19	Unknown	U36595†
b9, 10	<i>GBF-1</i>	G-box binding factor 1	X63894
b11, 12	EST23	Elongation factor	X52256
c1, 2	EST29	Aldolase	T04477
c3, 4	<i>GBF-2</i>	G-box binding factor 2	X63895
c5, 6	EST34	Chloroplast protease	R87034
c7, 8	EST35	Unknown	T14152
c9, 10	EST41	Catalase	T22720
c11, 12	rGR	Rat glucocorticoid receptor	M14053
d1, 2	EST42	Unknown	U36596†
d3, 4	EST45	ATPase	J04185
d5, 6	<i>HAT1</i>	Homeobox-leucine zipper 1	U09332
d7, 8	EST46	Light harvesting complex	T04063
d9, 10	EST49	Unknown	T76267
d11, 12	<i>HAT2</i>	Homeobox-leucine zipper 2	U09335
e1, 2	<i>HAT4</i>	Homeobox-leucine zipper 4	M90394
e3, 4	EST50	Phosphoribulokinase	T04344
e5, 6	<i>HAT5</i>	Homeobox-leucine zipper 5	M90416
e7, 8	EST51	Unknown	Z33675
e9, 10	<i>HAT22</i>	Homeobox-leucine zipper 22	U09336
e11, 12	EST52	Oxygen evolving	T21749
f1, 2	EST59	Unknown	Z34607
f3, 4	<i>KNAT1</i>	Knotted-like homeobox 1	U14174
f5, 6	EST60	RuBisCO small subunit	X14564
f7, 8	EST69	Translation elongation factor	T42799
f9, 10	<i>PPH1</i>	Protein phosphatase 1	U34803
f11, 12	EST70	Unknown	T44621
g1, 2	EST75	Chloroplast protease	T43698
g3, 4	EST78	Unknown	R65481
g5, 6	<i>ROC1</i>	Cyclophilin	L14844
g7, 8	EST82	GTP binding	X59152
g9, 10	EST83	Unknown	Z33795
g11, 12	EST84	Unknown	T45278
h1, 2	EST91	Unknown	T13832
h3, 4	EST96	Unknown	R64816
h5, 6	<i>SAR1</i>	Synaptobrevin	M90418
h7, 8	EST100	Light harvesting complex	Z18205
h9, 10	EST103	Light harvesting complex	X03909
h11, 12	<i>TRP4</i>	Yeast tryptophan biosynthesis	X04273

\*Proprietary sequence of Stratagene (La Jolla, California).

†No match in the database; novel EST.

REFERENCES AND NOTES

rent EST database (dbEST release 091495) e National Center for Biotechnology Information (Bethesda, MD) contains a total of 322,225 including 255,645 from the human genome 044 from Arabidopsis. Access is available via the Wide Web (<http://www.ncbi.nlm.nih.gov>).  
Meyerowitz and R. E. Pruitt, *Science* 229, 1214 (1986); R. E. Pruitt and E. M. Meyerowitz, *J. Mol. Biol.* 9 (1986); I. Hwang et al., *Plant J.* 1, 367 (1991); S. et al., *Plant Mol. Biol.* 24, 685 (1994); L. Le et al., *Mol. Gen. Genet.* 245, 390 (1994).  
on thesis, Stanford University (1995);  
O. Brown, in preparation. Microarrays were ed on poly-L-lysine-coated microscope Sigma) with a custom-built arraying machine ith one printing tip. The tip loaded 1 µl of PCR t (0.5 mg/ml) from 96-well microtiter plates osited -0.005 µl per slide on 40 slides at a y of 500 µm. The printed slides were rehydrat- 2 hours in a humid chamber, snap-dried at for 1 min, rinsed in 0.1% SDS, and treated 05% succinic anhydride prepared in buffer ng of 50% 1-methyl-2-pyrrolidone and nc acid. The cDNA on the slides was dena- distilled water for 2 min at 90°C immediately use. Microarrays were scanned with a laser ent scanner that contained a computer-con- Y stage and a microscope objective. A moded ultiline laser allowed sequential excitation of luorophores. Emitted light was split accord- ravelength and detected with two photomul- tes. Signals were read into a PC with the use -bit analog-to-digital board. Additional details array fabrication and use may be obtained by of e-mail ([pbrown@cmgm.stanford.edu](mailto:pbrown@cmgm.stanford.edu)).  
usubel et al., Eds., *Current Protocols in Mo- Biology* (Greene & Wiley Interscience, New 994), pp. 4.3.1-4.3.4.  
nylated [poly(A)<sup>+</sup>] mRNA was prepared from A with the use of Oligotex-dT resin (Qiagen). r transcription (RT) reactions were carried out itraScript RT-PCR kit (Stratagene) modified ws: 50-µl reactions contained 0.1 µg/µl of osis mRNA, 0.1 ng/µl of human AChR 0.05 µg/µl of oligo(dT) (21-mer), 1× first buffer, 0.03 U/µl of ribonuclease block, 500 xyadenosine triphosphate (dATP), 500 µM anosine triphosphate, 500 µM dTTP, 40 xyctosine triphosphate (dCTP), 40 µM flu- 12-dCTP (or isamine-5-dCTP), and 0.03 StrataScript reverse transcriptase. Reactions ubated for 60 min at 37°C, precipitated with , and resuspended in 10 µl of TE (10 mM tris- 1 mM EDTA, pH 8.0). Samples were then for 3 min at 94°C and chilled on ice. The RNA graded by adding 0.25 µl of 10 N NaOH 1 by a 10-min incubation at 37°C. The same- re neutralized by addition of 2.5 µl of 1 M H 8.0) and 0.25 µl of 10 N HCl and precip- ith ethanol. Pellets were washed with 70% dned to completion in a speedvac, resus- in 10 µl of H<sub>2</sub>O, and reduced to 3.0 µl in a ac. Fluorescent nucleotide analogs were ob- om New England Nuclear (DuPont).  
ation reactions contained 1.0 µl of fluorescent ythesis product (5) and 1.0 µl of hybridization 10× saline sodium citrate (SSC) and 0.2% e 2.0-µl probe mixtures were aliquoted onto array surface and covered with cover slips round). Arrays were transferred to a hybrid- chamber (3) and incubated for 18 hours at rays were washed for 5 min at room temper- 5°C) in low-stringency wash buffer (1× SSC % SDS), then for 10 min at room temperature stringency wash buffer (0.1× SSC and 0.1% rays were scanned in 0.1× SSC with the use rescence laser-scanning device (3).  
i of poly(A)<sup>+</sup> mRNA (4, 5) were spotted onto embranes (Nytan) and crosslinked with ul- ight with the use of a Stratalinker 1800 ene). Probes were prepared by random with the use of a Prime-It II kit (Stratagene) in ence of [<sup>32</sup>P]dATP. Hybridizations were car- according to the instructions of the manu-

facturer. Quantitation was performed on a Pho-  
phorimager (Molecular Dynamics).  
8. M. Schena and R. W. Davis, *Proc. Natl. Acad. Sci. U.S.A.* 89, 3894 (1992); M. Schena, A. M. Lloyd, R. W. Davis, *Genes Dev.* 7, 367 (1993); M. Schena and R. W. Davis, *Proc. Natl. Acad. Sci. U.S.A.* 91, 8393 (1994).  
9. H. Hottel et al., *Plant J.* 4, 1051 (1993); T. Newman et al., *Plant Physiol.* 106, 1241 (1994).  
10. N. E. Morton, *Proc. Natl. Acad. Sci. U.S.A.* 88, 7474 (1991); E. D. Green and R. H. Waterston, *J. Am. Med. Assoc.* 266, 1966 (1991); C. Betanhe-Chantelot, *Cell* 70, 1059 (1992); D. R. Cox et al., *Science* 265, 2031 (1994).  
11. E. S. Kawasaki et al., *Proc. Natl. Acad. Sci. U.S.A.* 85, 5698 (1988).

12. The laser fluorescent scanner was designed and fabri-  
cated in collaboration with S. Smith of Stanford Univer-  
sity. Scanner and analysis software was developed by  
R. X. Xia. The succinic anhydride reaction was sugges-  
ted by J. Muligan and J. Van Ness of Darwin Molecular  
Corporation. Thanks to S. Theologis, C. Somerville, K.  
Yamamoto, and members of the laboratories of R.W.C.  
and P.O.B. for critical comments. Supported by the  
Howard Hughes Medical Institute and by grants from  
NIH (R21HG00450) (P.O.B.) and R37AG00196  
(R.W.D.) and from NSF (MCB9106011) (R.W.D.) and  
by an NSF graduate fellowship (D.S.). P.O.B. is an  
assistant investigator of the Howard Hughes Medical  
Institute.

11 August 1995; accepted 22 September 1995

Gene Therapy in Peripheral Blood  
Lymphocytes and Bone Marrow for  
ADA<sup>-</sup> Immunodeficient Patients

Claudio Bordignon,\* Luigi D. Notarangelo, Nadia Nobili,  
Giuliana Ferrari, Giulia Casorati, Paola Panina, Evelina Mazzolari,  
Daniela Maggioni, Claudia Rossi, Paolo Servida,  
Alberto G. Ugazio, Fulvio Mavilio

Adenosine deaminase (ADA) deficiency results in severe combined immunodeficiency, the first genetic disorder treated by gene therapy. Two different retroviral vectors were used to transfer ex vivo the human ADA minigene into bone marrow cells and peripheral blood lymphocytes from two patients undergoing exogenous enzyme replacement therapy. After 2 years of treatment, long-term survival of T and B lymphocytes, marrow cells, and granulocytes expressing the transferred ADA gene was demonstrated and resulted in normalization of the immune repertoire and restoration of cellular and humoral immunity. After discontinuation of treatment, T lymphocytes, derived from transduced peripheral blood lymphocytes, were progressively replaced by marrow-derived T cells in both patients. These results indicate successful gene transfer into long-lasting progenitor cells, producing a functional multilineage progeny.

Severe combined immunodeficiency associated with inherited deficiency of ADA (1) is usually fatal unless affected children are kept in protective isolation or the immune system is reconstituted by bone marrow transplantation from a human leukocyte antigen (HLA)-identical sibling donor (2). This is the therapy of choice, although it is available only for a minority of patients. In recent years, other forms of therapy have been developed, including transplants from haploidentical donors (3, 4), exogenous enzyme replacement (5), and somatic-cell gene therapy (6-9).

We previously reported a preclinical model in which ADA gene transfer and expression

successfully restored immune functions in human ADA-deficient (ADA<sup>-</sup>) peripheral blood lymphocytes (PBLs) in immunodeficient mice in vivo (10, 11). On the basis of these preclinical results, the clinical application of gene therapy for the treatment of ADA<sup>-</sup> SCID (severe combined immunodeficiency disease) patients who previously failed exogenous enzyme replacement therapy was approved by our Institutional Ethical Committees and by the Italian National Committee for Bioethics (12). In addition to evaluating the safety and efficacy of the gene therapy procedure, the aim of the study was to define the relative role of PBLs and hematopoietic stem cells in the long-term reconstitution of immune functions after retroviral vector-mediated ADA gene transfer. For this purpose, two structurally identical vectors expressing the human ADA complementary DNA (cDNA), distinguishable by the presence of alternative restriction sites in a nonfunctional region of the viral long-terminal repeat (LTR), were used to transduce PBLs and bone marrow (BM) cells independently. This procedure allowed identification of the origin of

C. Bordignon, N. Nobili, G. Ferrari, D. Maggioni, C. Rossi, P. Servida, F. Mavilio, Telethon Gene Therapy Program for Genetic Diseases, DIBIT, Istituto Scientifico H. S. Raffaele, Milan, Italy.  
L. D. Notarangelo, E. Mazzolari, A. G. Ugazio, Department of Pediatrics, University of Brescia Medical School, Brescia, Italy.  
G. Casorati, Unità di Immunochimica, DIBIT, Istituto Scientifico H. S. Raffaele, Milan, Italy.  
P. Panina, Roche Milano Ricerche, Milan, Italy.

\*To whom correspondence should be addressed.

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**

**This Page Blank (uspto)**